



Types of Wounds

Skin Ulcers

Skin ulcers are round open sores on the skin caused by poor blood circulation. Skin ulcer treatments may include advanced dressings and wraps, antibiotics, pain management, and in rare cases, surgery.

Venous Ulcer

Venous ulcers occur when leg veins don't properly return blood back to the heart because of poor blood circulation in the legs. They usually form on the sides of the lower leg, above the ankles and below the calves. If left untreated, it can lead to infection or even blood poisoning.

Pressure Ulcer

A pressure ulcer is also known as a bed sore, are areas of damaged skin caused by unrelieved pressure. Pressure ulcers usually form around the ankles, back, elbows, heels, and hips.

Diabetic Ulcer

A diabetic ulcer is an open sore or wound that occurs people with diabetes, due to poor circulation, nerve damage, and high blood sugar.

Neuropathic Ulcer

A neuropathic ulcer is caused by repeated stress and pressure that diminishes sensation. These ulcers are typically seen in diabetes patients.

Arterial Ulcer

An arterial ulcer occurs when plaque builds up in the arteries reducing the amount of blood delivered to the body.

Other Wounds We Treat

- Gangrene- dead tissue caused by an infection or lack of blood flow
- Tissue and bone damage caused by radiation or infection
- Surgical wounds and post-op infections
- Skin irritations
- Osteoradionecrosis
- Bone infection (osteomyelitis)
- Other...

If you would like additional information or would like to schedule an appointment, please feel free to call the Center today: 661-634-0200.